

Tilburg University

The D1-triangulation in simplicial variable dimension algorithms for computing solutions of nonlinear equations

Dang, C.; Talman, A.J.J.

Publication date:
1990

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):

Dang, C., & Talman, A. J. J. (1990). *The D1-triangulation in simplicial variable dimension algorithms for computing solutions of nonlinear equations*. (CentER Discussion Paper; Vol. 1990-50). Unknown Publisher.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

CBM

entER

CBM

R R

8414

1990

for
mic Research

Discussion paper

50



No. 9050

THE D_1 -TRIANGULATION IN SIMPLICIAL VARIABLE
DIMENSION ALGORITHMS ON THE UNIT SIMPLEX
FOR COMPUTING FIXED POINTS

by Chuangyin Dang and Dolf Talman *R 83*

515.14
696.42

August 1990

ISSN 0924-7815

The D_1 -Triangulation in Simplicial Variable Dimension Algorithms on the Unit Simplex for Computing Fixed Points

Chuangyin Dang and Dolf Talman
CentER for Economic Research
Tilburg University
The Netherlands

Abstract

We consider how to use the D_1 -triangulation in simplicial variable dimension algorithms on the unit simplex for computing fixed points. A new version of the D_1 -triangulation is proposed. We can use directly this version in simplicial variable dimension algorithms on the unit simplex. According to measures of efficiency of triangulations, the D_1 -triangulation is superior to the other well-known triangulations of R^n . So it is hoped that the cost of computation can be reduced through using the D_1 -triangulation in simplicial variable dimension algorithms on the unit simplex.

Keywords: Simplicial Variable Dimension Algorithms, Triangulations, Measures of Efficiency of Triangulations

1 Introduction

In order to find out fixed points of nonlinear equations, a number of simplicial algorithms have been introduced. A simplicial algorithm subdivides the underlying space into simplices and searches for a simplex that yields an approximate solution. To find such a simplex, the algorithm generates a sequence of adjacent simplices. A simplicial variable dimension algorithm generates a sequence of simplices of varying dimension. This sequence connects an arbitrary starting point with an approximate solution. Since van der Laan and Talman proposed the first simplicial variable dimension algorithm without an extra dimension on the unit simplex $S^n = \{x \in R_+^{n+1} \mid \sum_{j=0}^n x_j = 1\}$ in [8], several contributions have been made, for example, their $(n+1)$ -ray and $2n$ -ray methods on R^n in [9], Wright's 2^n -ray method on R^n in [13], Kojima and Yamamoto's $(3^n - 1)$ -ray method on R^n in [7], Yamamoto's 2-ray method on R^n in [14], Doup, van der Laan and Talman's $(2^{n+1} - 2)$ -ray method on S^n in [5], and so on. For a survey on simplicial algorithms, see [1]. In [10], a simplicial variable dimension algorithm was introduced by Talman and Yamamoto to find a stationary point of a continuous function on a polytope. Doup gave an excellent survey about simplicial variable dimension algorithms on the product space of unit simplices in [4]. Recently, a new triangulation of R^n was proposed in [2], the D_1 -triangulation. According to measures of efficiency of triangulations, it is superior to the other well-known triangulations. But it is not straightforward to use the D_1 -triangulation in simplicial variable dimension algorithms except the $2n$ -ray and 2-ray methods on R^n . In [3], we introduced a version of the D_1 -triangulation for simplicial variable dimension algorithms on the unit simplex. But it has as shortcoming not to be able to induce a simplicial subdivision according to the D_1 -triangulation of all the subsets into which a simplicial variable dimension algorithm subdivides the unit simplex. To reduce the cost of computation, we reconsider how to apply the D_1 -triangulation to simplicial variable dimension algorithms on the unit simplex. We propose a new version of the D_1 -triangulation, called the D_{v1} -triangulation. This version subdivides each of the subsets according to the D_1 -triangulation into which a simplicial algorithm subdivides the unit simplex.

The second section introduces the D_1 -triangulation. The third section gives the pivot rules of the D_{v1} -triangulation. The fourth section presents

how to apply the D_{v1} -triangulation to simplicial variable dimension algorithms on the unit simplex.

2 The D_{v1} -Triangulation

In this section we describe the D_{v1} -triangulation. This simplicial subdivision is based on the D_1 -triangulation of R^n introduced in [2]. The D_{v1} -triangulation will be used as the underlying subdivision for simplicial variable dimension algorithms on the unit simplex. An n -dimensional simplex is the convex hull of $n + 1$ affinely independent points called its vertices, and a simplicial subdivision of a compact, convex set is the set of a finite number of simplices such that the intersection of each pair of simplices is either empty or a common face and such that the union of all simplices is the set itself.

Let m be a positive integer and let N denote the index set $\{1, 2, \dots, n\}$. Furthermore, let

$$C^n(m) = \{x \in R^n \mid m \geq x_1 \geq x_2 \geq \dots \geq x_n \geq 0\}.$$

The D_{v1} -triangulation subdivides simplicially the set $C^n(m)$. The simplices of this triangulation can be represented by some specific vector y , permutation π , sign vector s , and integer p . Let D denote the set

$$\{y \in C^n(m) \mid \text{all components of } y \text{ are even}\}.$$

Take any $y \in D$. Let $1 \leq h \leq n$ be such that for $j = 1, 2, \dots, h$, it holds that $y_{m_j+1} = y_{m_j+k}$ for $k = 2, 3, \dots, n_j$, and for $j = 1, 2, \dots, h - 1$, it holds that $y_{m_j+n_j} > y_{m_{j+1}+1}$, where $m_1 = 0$, $n_h = n - m_h$, and $m_j + n_j = m_{j+1}$ for $j = 1, 2, \dots, h - 1$.

Take $s = (s_1, s_2, \dots, s_n)^T$ to be a sign vector such that

$$s_{m_j+1} = \dots = s_{m_j+k_j} = 1$$

and

$$s_{m_j+k_j+1} = \dots = s_{m_j+n_j} = -1$$

for $j = 1, 2, \dots, h$, where $k_1 = 0$ if $y_1 = m$, $k_h = n_h$ if $y_n = 0$, and $0 \leq k_j \leq n_j$ for $j = 1, 2, \dots, h$.

For $1 \leq j \leq h$ and $1 \leq k \leq k_j$, let n -vector $g(m_j + k)$ be given by

$$g_i(m_j + k) = \begin{cases} 1 & \text{if } m_j + 1 \leq i \leq m_j + k, \\ 0 & \text{otherwise,} \end{cases}$$

for $i = 1, 2, \dots, n$.

For $1 \leq j \leq h$ and $k_j + 1 \leq k \leq n_j$, let n -vector $g(m_j + k)$ be given by

$$g_i(m_j + k) = \begin{cases} -1 & \text{if } m_j + k \leq i \leq m_j + n_j, \\ 0 & \text{otherwise,} \end{cases}$$

for $i = 1, 2, \dots, n$.

Take an integer p such that $0 \leq p \leq n - 1$ and if $h = 1$ and $k_1 = 0$ or n then $p = 0$.

Finally, take a permutation π of the elements of N such that for $1 \leq j \leq h$,

$$\pi^{-1}(m_j + 1) > \pi^{-1}(m_j + 2) > \dots > \pi^{-1}(m_j + k_j)$$

and

$$\pi^{-1}(m_j + k_j + 1) < \pi^{-1}(m_j + k_j + 2) < \dots < \pi^{-1}(m_j + n_j)$$

and when $p \geq 1$, there exists no $1 \leq j \leq h$ such that $m_j + 1 \leq \pi(k) \leq m_j + k_j$ for all $p \leq k \leq n$ or $m_j + k_j + 1 \leq \pi(k) \leq m_j + n_j$ for all $p \leq k \leq n$.

Let u^i denote the i -th unit vector in \mathbb{R}^n for $i = 1, 2, \dots, n$.

Definition 2.1. Let the vector y , the permutation π , the sign vector s , and the number p be taken as above. Then the vectors y^0, y^1, \dots, y^n are given as follows.

$$\begin{aligned} &\text{If } p = 0, \text{ then } y^0 = y, \\ &y^k = y + g(\pi(k)), k = 1, 2, \dots, n. \\ &\text{If } p \geq 1, \text{ then } y^0 = y + s, \\ &y^k = y^{k-1} - s_{\pi(k)} u^{\pi(k)}, k = 1, 2, \dots, p - 1, \\ &y^k = y + g(\pi(k)), k = p, p + 1, \dots, n. \end{aligned}$$

Let y^0, y^1, \dots, y^n be obtained from the above definition. Then it is obvious that they are affinely independent. Thus their convex hull is a simplex with vertices y^0, y^1, \dots, y^n . Let us denote this simplex by $D_{v1}(y, \pi, s, p)$. Let D_{v1} be the set of all such simplices $D_{v1}(y, \pi, s, p)$. We will show that D_{v1} is a simplicial subdivision of $C^n(m)$.

For given y and s as above, let α be defined by $\alpha = \sum_{j=1}^h \alpha_j$, where

$$\alpha_j = \begin{cases} 2 & \text{if } 0 < k_j < n_j, \\ 1 & \text{otherwise,} \end{cases}$$

for $j = 1, 2, \dots, h$.

For given y, π, s , and p as above, let

$$r_j = | \{m_j + k \mid 1 \leq k \leq k_j\} \cap \{\pi(k) \mid 1 \leq k \leq p-1\} |$$

and

$$q_j = | \{m_j + k \mid k_j + 1 \leq k \leq n_j\} \cap \{\pi(k) \mid 1 \leq k \leq p-1\} |$$

for $j = 1, 2, \dots, h$.

Lemma 2.2. The union of all $\sigma \in D_{v1}$ is equal to $C^n(m)$.

Proof. Clearly, every $\sigma \in D_{v1}$ is a subset of $C^n(m)$. Let $x \in C^n(m)$ be arbitrary. Then $x \in D_{v1}(y, \pi, s, p)$ with y, π, s , and p determined as follows.

Take the vector y equal to

$$y_i = \begin{cases} \lfloor x_i \rfloor + 1 & \text{if } \lfloor x_i \rfloor \text{ is odd and } \lfloor x_i \rfloor < m, \\ \lfloor x_i \rfloor - 1 & \text{if } \lfloor x_i \rfloor \text{ is odd and } \lfloor x_i \rfloor = m, \\ \lfloor x_i \rfloor & \text{otherwise,} \end{cases}$$

for $i = 1, 2, \dots, n$ and the sign vector s equal to

$$s_i = \begin{cases} -1 & \text{if } \lfloor x_i \rfloor \text{ is odd and } \lfloor x_i \rfloor < m \\ & \text{or } \lfloor x_i \rfloor \text{ is even and } \lfloor x_i \rfloor = m, \\ 1 & \text{otherwise,} \end{cases}$$

for $i = 1, 2, \dots, n$. It is obvious that $y \in D$. Take the permutation π such that

$$s_{\pi(1)}(x_{\pi(1)} - y_{\pi(1)}) \leq s_{\pi(2)}(x_{\pi(2)} - y_{\pi(2)}) \leq \dots \leq s_{\pi(n)}(x_{\pi(n)} - y_{\pi(n)})$$

and for $1 \leq j \leq h$,

$$\pi^{-1}(m_j + 1) > \pi^{-1}(m_j + 2) > \dots > \pi^{-1}(m_j + k_j)$$

and

$$\pi^{-1}(m_j + k_j + 1) < \pi^{-1}(m_j + k_j + 2) < \cdots < \pi^{-1}(m_j + n_j).$$

Suppose $\alpha = 1$. Then take $p = 0$. Now, let $y^0 = y$,

$$y^k = y + g(\pi(k)), k = 1, 2, \dots, n,$$

and if $s_1 = 1$, let

$$\begin{aligned}\beta_0 &= 1 - s_1(x_1 - y_1), \\ \beta_1 &= s_n(x_n - y_n), \\ \beta_2 &= s_{n-1}(x_{n-1} - y_{n-1}) - s_n(x_n - y_n), \\ &\dots \\ \beta_n &= s_1(x_1 - y_1) - s_2(x_2 - y_2),\end{aligned}$$

and if $s_1 = -1$, let

$$\begin{aligned}\beta_0 &= 1 - s_n(x_n - y_n), \\ \beta_1 &= s_1(x_1 - y_1), \\ \beta_2 &= s_2(x_2 - y_2) - s_1(x_1 - y_1), \\ &\dots \\ \beta_n &= s_n(x_n - y_n) - s_{n-1}(x_{n-1} - y_{n-1}).\end{aligned}$$

Obviously, $\sum_{k=0}^n \beta_k = 1, \beta_k \geq 0$ for all k , and

$$x = \sum_{k=0}^n \beta_k y^k.$$

Hence,

$$x \in D_{v1}(y, \pi, s, p).$$

When $\alpha = n$, the proof is the same as that of **Lemma 2.2** in [2].

Now suppose that $1 < \alpha < n$. For $1 \leq j \leq h$, let

$$\mu_j = \begin{cases} s_{m_j+1}(x_{m_j+1} - y_{m_j+1}) + s_{m_j+n_j}(x_{m_j+n_j} - y_{m_j+n_j}) & \text{if } \alpha_j = 2, \\ s_{m_j+1}(x_{m_j+1} - y_{m_j+1}) & \text{if } \alpha_j = 1 \text{ and } s_{m_j+1} = 1, \\ s_{m_j+n_j}(x_{m_j+n_j} - y_{m_j+n_j}) & \text{if } \alpha_j = 1 \text{ and } s_{m_j+1} = -1. \end{cases}$$

Let $\mu = \sum_{j=1}^h \mu_j$.

In case $\mu \leq 1$. Take $p = 0$, and let $y^0 = y$,

$$y^k = y + g(\pi(k)), k = 1, 2, \dots, n.$$

For $1 \leq k \leq n$, let

$$\beta_k = \begin{cases} s_{\pi(k)}(x_{\pi(k)} - y_{\pi(k)}) & \text{if } \pi(k) = m_j + k_j \text{ and } k_j \geq 1 \text{ or } \pi(k) = \\ & m_j + k_j + 1 \text{ and } k_j \leq n_j - 1 \text{ for some } 1 \leq j \leq h, \\ s_{\pi(k)}(x_{\pi(k)} - y_{\pi(k)}) - s_{\pi(k)+1}(x_{\pi(k)+1} - y_{\pi(k)+1}) & \text{if } 1 \leq \pi(k) - m_j < k_j \text{ for some } 1 \leq j \leq h, \\ s_{\pi(k)}(x_{\pi(k)} - y_{\pi(k)}) - s_{\pi(k)-1}(x_{\pi(k)-1} - y_{\pi(k)-1}) & \text{if } k_j + 1 < \pi(k) - m_j \leq n_j \text{ for some } 1 \leq j \leq h, \end{cases}$$

and let $\beta_0 = 1 - \mu$. Then it is clear that $\beta_k \geq 0$ for all k , $\sum_{k=0}^n \beta_k = 1$, and

$$x = \sum_{k=0}^n \beta_k y^k.$$

Thus

$$x \in D_{v1}(y, \pi, s, p).$$

Now suppose $\mu > 1$. Let p_{max} denote the largest $1 \leq p \leq n-1$ such that there exists no $1 \leq j \leq h$ such that $m_j + 1 \leq \pi(k) \leq m_j + k_j$ for all $p \leq k \leq n$ or $m_j + k_j + 1 \leq \pi(k) \leq m_j + n_j$ for all $p \leq k \leq n$. We show that there exists $1 \leq p \leq p_{max}$ such that the following system has a nonnegative solution,

$$\begin{aligned} \beta_0 &= s_{\pi(1)}(x_{\pi(1)} - y_{\pi(1)}), \\ \beta_1 &= s_{\pi(2)}(x_{\pi(2)} - y_{\pi(2)}) - s_{\pi(1)}(x_{\pi(1)} - y_{\pi(1)}), \\ &\dots \\ \beta_{p-2} &= s_{\pi(p-1)}(x_{\pi(p-1)} - y_{\pi(p-1)}) \\ &\quad - s_{\pi(p-2)}(x_{\pi(p-2)} - y_{\pi(p-2)}), \\ \beta_{p-1} &= -s_{\pi(p-1)}(x_{\pi(p-1)} - y_{\pi(p-1)}) \\ &\quad + (\sum_{i=p}^n \lambda_{\pi(i)} - 1)/(c(p) - 1), \end{aligned}$$

$$\beta_k = \begin{cases} s_{\pi(k)}(x_{\pi(k)} - y_{\pi(k)}) - s_{\pi(k)+1}(x_{\pi(k)+1} - y_{\pi(k)+1}) & \text{if } 1 \leq \pi(k) - m_j < k_j - r_j \text{ for some } 1 \leq j \leq h, \\ s_{\pi(k)}(x_{\pi(k)} - y_{\pi(k)}) - (\sum_{i=p}^n \lambda_{\pi(i)} - 1)/(c(p) - 1) & \text{if } 1 \leq \pi(k) - m_j = k_j - r_j \\ & \text{or } k_j + q_j + 1 = \pi(k) - m_j \leq n_j \text{ for some } 1 \leq j \leq h, \\ s_{\pi(k)}(x_{\pi(k)} - y_{\pi(k)}) - s_{\pi(k)-1}(x_{\pi(k)-1} - y_{\pi(k)-1}) & \text{if } k_j + q_j + 1 < \pi(k) - m_j \leq n_j \text{ for some } 1 \leq j \leq h, \end{cases}$$

for $k = p, p+1, \dots, n$, where for $i = p, \dots, n$,

$$\lambda_{\pi(i)} = \begin{cases} 0 & \text{if } 1 < \pi(i) - m_j \leq k_j - r_j \\ & \text{or } k_j + q_j + 1 \leq \pi(i) - m_j < n_j \\ & \text{for some } 1 \leq j \leq h, \\ s_{m_j+1}(x_{m_j+1} - y_{m_j+1}) & \text{if } 1 = \pi(i) - m_j \leq k_j - r_j \\ & \text{for some } 1 \leq j \leq h, \\ s_{m_j+n_j}(x_{m_j+n_j} - y_{m_j+n_j}) & \text{if } k_j + q_j + 1 \leq \pi(i) - m_j = n_j \\ & \text{for some } 1 \leq j \leq h, \end{cases}$$

and

$$c(p) = \sum_{j=1}^h c_j(p)$$

with

$$c_j(p) = \begin{cases} 0 & \text{if } r_j = k_j \text{ and } q_j = n_j - k_j, \\ 2 & \text{if } r_j < k_j \text{ and } q_j < n_j - k_j, \\ 1 & \text{otherwise,} \end{cases}$$

for $j = 1, 2, \dots, h$.

If $\beta_{p-1} \geq 0$ for $p = p_{max}$, it is clear that $\beta_k \geq 0$ for all k . If not, then since $\mu > 1$, there exists $1 \leq p_0 \leq p_{max} - 1$ such that

$$0 \leq -s_{\pi(p_0-1)}(x_{\pi(p_0-1)} - y_{\pi(p_0-1)}) + (\sum_{i=p_0}^n \lambda_{\pi(i)} - 1)/(c(p_0) - 1)$$

and

$$0 > -s_{\pi(p_0)}(x_{\pi(p_0)} - y_{\pi(p_0)}) + (\sum_{i=p_0+1}^n \lambda_{\pi(i)} - 1)/(c(p_0 + 1) - 1).$$

Hence,

$$s_{\pi(p_0)}(x_{\pi(p_0)} - y_{\pi(p_0)}) - (\sum_{i=p_0}^n \lambda_{\pi(i)} - 1)/(c(p_0) - 1) \geq 0.$$




FIGURE 1

Table 1: The Pivot Rules of the D_v -Triangulation

i	p			y	s	π	\bar{p}
0	0	$\alpha = 1, s_1 = 1$	$y_1 = m - 1$	BD(1)			
			otherwise	$y + 2s_1 u^1$	$s - 2s_1 u^1$	π	$n - 1$
		$\alpha = 1, s_1 = -1$		$y + 2s_n u^n$	$s - 2s_n u^n$	π	$n - 1$
0	1	$\alpha \geq 2$		y	s	π	$p + 1$
				y	s	π	$p - 1$
0	$p \geq 2$		$y_{\pi(1)} = 0$ or m	BD(2)			
			otherwise	y	$s - 2s_{\pi(1)} u^{\pi(1)}$	π	p
$1 \leq i \leq n$	0	$1 \leq \pi(i) - m_j = k_j$ or $k_j + 1 = \pi(i) - m_j \leq n_j$ for some $1 \leq j \leq h$	$y_{\pi(i)} = 0$ or m	BD(3)			
		otherwise	otherwise	y	$s - 2s_{\pi(i)} u^{\pi(i)}$	π	p
$1 \leq i < p - 1$		otherwise		BD(4)			
		for some $1 \leq j \leq h$, $1 \leq \pi(k) - m_j \leq k_j$ for $k = i, i + 1$ or $k_j + 1 \leq \pi(k) - m_j \leq n_j$ for $k = i, i + 1$		BD(5)			
		otherwise		y	s	$(\pi(1), \dots, \pi(i + 1), \pi(i), \dots, \pi(n))$	p
$i = p - 1$	$p \geq 2$			y	s	π	$p - 1$
$p \leq i \leq n$	$1 \leq p < n - 1$	for some $1 \leq j \leq h$, $1 \leq \pi(k) - m_j \leq k_j$ for $p \leq k \leq n$ and $k \neq i$ or $k_j + 1 \leq \pi(k) - m_j \leq n_j$ for $p \leq k \leq n$ and $k \neq i$	$i < n$ and either $y_{\pi(n)} \neq m - 1$ or $s_{\pi(n)} \neq 1$	$y + 2s_{\pi(n)} u^{\pi(n)}$	$s - 2s_{\pi(n)} u^{\pi(n)}$	$(\pi(1), \dots, \pi(p - 1), \pi(i), \pi(p), \dots, \pi(i - 1), \pi(i + 1), \dots, \pi(n))$	$n - 1$
			$i = n$ and either $y_{\pi(n-1)} \neq m - 1$ or $s_{\pi(n-1)} \neq 1$	$y + 2s_{\pi(n-1)} u^{\pi(n-1)}$	$s - 2s_{\pi(n-1)} u^{\pi(n-1)}$	$(\pi(1), \dots, \pi(p - 1), \pi(n), \pi(p), \dots, \pi(n - 1))$	$n - 1$
		for some $1 \leq j \leq h$, $1 \leq \pi(i) - m_j = k_j - r_j$ or $k_j + q_j + 1 = \pi(i) - m_j \leq n_j$	otherwise	BD(6)			
				y	s	$(\pi(1), \dots, \pi(p - 1), \pi(i), \pi(p), \dots, \pi(i - 1), \pi(i + 1), \dots, \pi(n))$	$p + 1$
	$1 \leq p = n - 1$	otherwise		BD(7)			
			$i = n - 1$ and either $y_{\pi(n)} \neq m - 1$ or $s_{\pi(n)} \neq 1$	$y + 2s_{\pi(n)} u^{\pi(n)}$	$s - 2s_{\pi(n)} u^{\pi(n)}$	π	p^*
		otherwise	$i = n$ and either $y_{\pi(n-1)} \neq m - 1$ or $s_{\pi(n-1)} \neq 1$	$y + 2s_{\pi(n-1)} u^{\pi(n-1)}$	$s - 2s_{\pi(n-1)} u^{\pi(n-1)}$	π^*	p^*
			otherwise	BD(8)			

Thus when $p = p_0$, $\beta_k \geq 0$ for all k . Obviously, $\sum_{k=0}^n \beta_k = 1$. Let $y^0 = y + s$, and for $k = 1, 2, \dots, p-1$, let

$$y^k = y^{k-1} - s_{\pi(k)} u^{\pi(k)},$$

and for $k = p, p+1, \dots, n$, let

$$y^k = y + g(\pi(k)).$$

We easily obtain that $x = \sum_{k=0}^n \beta_k y^k$. Therefore,

$$x \in D_{v1}(y, \pi, s, p).$$

From the above conclusions, the lemma follows immediately.

Theorem 2.3. D_{v1} is a triangulation of $C^n(m)$.

Proof. From **Lemma 2.2** and the definition, the theorem follows immediately.

We call this simplicial subdivision of $C^n(m)$ the D_{v1} -triangulation. The D_{v1} -triangulation of $C^3(2)$ is illustrated in Figure 1. The D_{v1} -triangulation of $C^n(m)$ is such that the set $C^n(m)$ is subdivided into simplices according to the D_1 -triangulation of R^n with grid size m^{-1} except the corners of the set. Moreover, every t -dimension face of $C^n(m)$ is also simplicially subdivided according to the D_1 -triangulation of R^t , $1 \leq t \leq n-1$. Therefore, the D_{v1} -triangulation of $C^n(m)$ is superior to all other triangulations of this set, see [2].

3 The Pivot Rules of the D_{v1} -Triangulation

Let $\sigma = D_{v1}(y, \pi, s, p)$ be a simplex of the D_{v1} -triangulation of $C^n(m)$ with vertices y^0, y^1, \dots, y^n . We want to obtain the parameters of the simplex $\bar{\sigma} = D_{v1}(\bar{y}, \bar{\pi}, \bar{s}, \bar{p})$ sharing with σ the facet opposite to the vertex y^i , $i \in \{0, 1, \dots, n\}$, of σ , in case this facet of σ does not lie in the boundary of $C^n(m)$. In Table 1, we show how $\bar{y}, \bar{\pi}, \bar{s}$ and \bar{p} depend on y, π, s, p and i . From this table, it is easy to obtain each vertex of $\bar{\sigma}$, and in particular the vertex opposite to the facet shared with σ .

In this table, π^* and p^* denote that if, for some $1 \leq j \leq \bar{h}$,

$$1 \leq \pi(k) - \bar{m}_j \leq \bar{k}_j \text{ for } k = n-1, n$$

or

$$\bar{k}_j + 1 \leq \pi(k) - \bar{m}_j \leq \bar{n}_j \text{ for } k = n-1, n,$$

then

$$\pi^* = (\pi(1), \dots, \pi(n-2), \pi(n), \pi(n-1))$$

and

$$p^* = \begin{cases} k & \text{if there exists } 1 \leq k \leq n-2 \text{ such that } 1 \leq \pi(l) - \bar{m}_j \leq \bar{k}_j \\ & \text{for all } k < l \leq n \text{ and } \pi(k) \text{ doesn't or} \\ & \bar{k}_j + 1 \leq \pi(l) - \bar{m}_j \leq \bar{n}_j \\ & \text{for all } k < l \leq n \text{ and } \pi(k) \text{ doesn't,} \\ 0 & \text{otherwise;} \end{cases}$$

otherwise, $\pi^* = \pi$ and $p^* = p$, where \bar{h} , \bar{m}_j , \bar{k}_j , and \bar{n}_j satisfy that for $j = 1, 2, \dots, \bar{h}$, it holds that $\bar{y}_{\bar{m}_j+1} = \bar{y}_{\bar{m}_j+k}$ for $k = 2, 3, \dots, \bar{n}_j$, and for $j = 1, 2, \dots, \bar{h}-1$, it holds that $\bar{y}_{\bar{m}_j+\bar{n}_j} > \bar{y}_{\bar{m}_{j+1}+1}$, where $\bar{m}_1 = 0$, $\bar{n}_{\bar{h}} = n - \bar{m}_{\bar{h}}$, and $\bar{m}_j + \bar{n}_j = \bar{m}_{j+1}$ for $j = 1, 2, \dots, \bar{h}-1$, and for $j = 1, 2, \dots, \bar{h}$, it holds that $\bar{s}_{\bar{m}_j+1} = \dots = \bar{s}_{\bar{m}_j+\bar{k}_j} = 1$ and $\bar{s}_{\bar{m}_j+\bar{k}_j+1} = \dots = \bar{s}_{\bar{m}_j+\bar{n}_j} = -1$, where $\bar{k}_1 = 0$ if $\bar{y}_1 = m$, and $\bar{k}_{\bar{h}} = \bar{n}_{\bar{h}}$ if $\bar{y}_{\bar{n}} = 0$.

4 Simplicial Variable Dimension Algorithms Based on the D_{v1} -Triangulation

Here we only consider how to use the D_{v1} -triangulation in the $(2^{n+1} - 2)$ -ray method proposed by Doup, van der Laan and Talman in [5]. It can similarly be derived how to use the D_{v1} -triangulation in the other simplicial variable dimension algorithms on the unit simplex.

Let N_0 denote the index set $\{0, 1, \dots, n\}$ and let S^n denote the unit simplex

$$\left\{ x \in R^{n+1} \mid \sum_{j=0}^n x_j = 1, x_j \geq 0 \text{ for all } j \in N_0 \right\}.$$

Let $f : S^n \rightarrow R^{n+1}$ be continuous on S^n such that $x^\top f(x) = 0$ for all $x \in S^n$, i.e., f is a complementary function on S^n . Our purpose is to find a point x^*

in S^n such that $f(x^*) \leq 0$. Take arbitrarily $x^0 \in S^n$ as an initial point. Let H be a nonempty subset of N_0 and let

$$S^n(H) = \{x \in S^n \mid x_j = 0 \text{ for all } j \notin H\}.$$

Set $H_0 = \{j \in H \mid x_j^0 = 0\}$. Then the projection vector of x^0 on $S^n(H)$, $v(H)$, is defined by

$$v_i(H) = \begin{cases} 0 & \text{if } i \notin H, \\ (1 - \sum_{j \in H} x_j^0) / (\sum_{j \in H} x_j^0 + |H_0|) & \text{if } i \in H_0, \\ x_i^0(1 + |H_0|) / (\sum_{j \in H} x_j^0 + |H_0|) & \text{otherwise,} \end{cases}$$

for $i = 0, 1, \dots, n$. When H is empty, we define $v(H) = x^0$.

Let $g = (g_0, g_1, \dots, g_n)^\top$ be a sign vector such that $g_i \in \{-1, 0, +1\}$ for $i = 0, 1, \dots, n$. Let $I^-(g) = \{i \in N_0 \mid g_i = -1\}$, $I^0(g) = \{i \in N_0 \mid g_i = 0\}$ and $I^+(g) = \{i \in N_0 \mid g_i = +1\}$. For a sign vector g such that both $I^+(g)$ and $\{i \in I^-(g) \mid x_i^0 > 0\}$ are nonempty, we define

$$A(g) = \left\{ x \in S^n \mid \begin{cases} x_k/x_k^0 = 1 + \alpha \text{ if } k \in I^+(g) \text{ and } x_k^0 > 0, \\ \beta \leq x_k/x_k^0 \leq 1 + \alpha \text{ if } k \in I^0(g) \text{ and } x_k^0 > 0, \\ \beta = x_k/x_k^0 \text{ if } k \in I^-(g) \text{ and } x_k^0 > 0, \\ x_k = \alpha \text{ if } k \in I^+(g) \text{ and } x_k^0 = 0, \\ 0 \leq x_k \leq \alpha \text{ if } k \in I^0(g) \text{ and } x_k^0 = 0, \\ 0 = x_k \text{ if } k \in I^-(g) \text{ and } x_k^0 = 0, \text{ and} \\ 0 \leq \beta \leq 1 \leq 1 + \alpha \end{cases} \right\}.$$

It is obvious that $A(g)$ is a t -dimensional polyhedron with $t = |I^0(g)| + 1$ and that S^n is equal to the union of $A(g)$ over all sign vectors g such that $|I^0(g)| = n - 1$. In addition, let $k_0 = 0$ and let $\gamma(g) = (k_1, k_2, \dots, k_{t-1})$ be a permutation of the $t - 1$ elements in $I^0(g)$. Then we define

$$A(g, \gamma(g)) = \left\{ x \in S^n \mid \begin{cases} x = x^0 + \sum_{j=0}^{t-1} \lambda(k_j) w(k_j) \\ \text{with } 0 \leq \lambda(k_{t-1}) \leq \dots \leq \lambda(k_0) \leq 1 \end{cases} \right\},$$

where $w(k_0) = v(I^+(g) \cup \{k_1, \dots, k_{t-1}\}) - x^0$ and

$$w(k_j) = v(I^+(g) \cup \{k_1, \dots, k_{t-j-1}\}) - v(I^+(g) \cup \{k_1, \dots, k_{t-j}\})$$

for $j = 1, 2, \dots, t - 1$. Then it is clear that $A(g, \gamma(g))$ is t -dimensional and that $A(g)$ is the union of $A(g, \gamma(g))$ over all permutations $\gamma(g)$ of the elements

in $I^0(g)$. Let P be the $(n+1) \times t$ -matrix with the i -th column equal to $w(k_{i-1})$ for $i = 1, 2, \dots, t$. Then

$$A(g, \gamma(g)) = \{x^0 + m^{-1}Px \mid x \in C^t(m)\}.$$

This means that $A(g, \gamma(g))$ is homeomorphic to $C^t(m)$. Thus the D_{v1} -triangulation induces a simplicial subdivision of $A(g, \gamma(g))$, which is denoted by $D_{v1}(g, \gamma(g))$. Moreover, as can easily be shown, the union of $D_{v1}(g, \gamma(g))$ over all permutations of the elements in $I^0(g)$ is a simplicial subdivision of $A(g)$. We write it as $D_{v1}(g)$. Finally, a simplicial subdivision of S^n has been obtained by taking the union of $D_{v1}(g)$ over all sign vectors g such that both $I^+(g)$ and $\{i \in I^-(g) \mid x_i^0 > 0\}$ are nonempty.

For $x \in R^{n+1}$, let the sign vector of x be defined by

$$\text{sgn}(x) = (\text{sgn}_0(x), \text{sgn}_1(x), \dots, \text{sgn}_n(x))^T,$$

where

$$\text{sgn}_i(x) = \begin{cases} -1 & \text{if } x_i < 0, \\ 0 & \text{if } x_i = 0, \\ +1 & \text{if } x_i = +1, \end{cases}$$

for $i = 0, 1, \dots, n$. Let $e(j)$ denote the j -th unit vector in R^{n+1} for $j = 0, 1, \dots, n$. For a given sign vector g with $t = |I^0(g)| + 1$, a k -simplex σ with vertices z^0, z^1, \dots, z^k , for $k = t - 1$ and t , is called g -complete if the $(n+2)$ -system of linear equations

$$\sum_{j=0}^k \lambda_j \begin{pmatrix} f(z^j) \\ 1 \end{pmatrix} - \sum_{j \notin I^0(g)} \mu_j g_j \begin{pmatrix} e(j) \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

has a nonnegative solution.

Starting at x^0 with $g = \text{sgn}(f(x^0))$, the $(2^{n+1} - 2)$ -ray method generates a sequence of adjacent g -complete t -simplices in $A(g)$ with g -complete common facets for varying sign vectors g . Below we describe the steps of the algorithm in case the simplicial subdivision of each set $A(g)$ is based on the D_{v1} -triangulation for some given grid size m^{-1} .

The $(2^{n+1} - 2)$ -Ray Method Based on the D_{v1} -Triangulation:

Initialization: Without loss of generality we assume that $f_i(x^0) \neq 0$ for all

i. Set $g = \text{sgn}(f(x^0))$ and $t = 1$. Set $y = 0$, $\pi = (1)$, $s = 1$ and $p = 0$. Further, set $z^0 = x^0$ and $\tau_0 = \{z^0\}$. Finally, set $k = 0$.

Step 1: Let σ_k be the simplex in $D_{v1}(g)$ corresponding to the simplex

$$D_{v1}(y, \pi, s, p).$$

Thus τ_k is a facet of σ_k . Let z^+ denote the vertex of σ_k opposite to τ_k . Perform a linear programming step with $l^+ = (f(z^+), 1)^\top$ in the system of linear equations

$$\sum_{j=0}^{t-1} \lambda_j \begin{pmatrix} f(z^j) \\ 1 \end{pmatrix} - \sum_{j \notin I^0(g)} \mu_j g_j \begin{pmatrix} e(j) \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

If some μ_c becomes zero, then set $z^t = z^+$ and go to **Step 2**; otherwise, some λ_d becomes zero, then set $z^- = z^d$ and $z^d = z^+$, and go to **Step 3**.

Step 2: When $I^+(g) = \{c\}$ or $\{j \in I^-(g) \mid x_j^0 > 0\} = \{c\}$, σ_k yields an approximate solution and the algorithm terminates; otherwise perform the following increasing dimension procedure. Set $\tau_{k+1} = \sigma_k$ and $k = k + 1$ and set $g_c = 0$. When $c \in I^+(g)$, set $\gamma(g) = (c, k_1, \dots, k_{t-1})$, $y_{t+1} = 0$ and $s_{t+1} = 1$ and if $p = 0$ then set $\pi = (t + 1, \pi(1), \dots, \pi(t))$ and if $p \geq 1$ then set $\pi = (t + 1, \pi(1), \dots, \pi(t))$ and $p = p + 1$. When $c \in I^-(g)$, set $\gamma(g) = (k_1, \dots, k_{t-1}, c)$, $y_i = y_{i-1}$ and $s_i = s_{i-1}$ for $i = t + 1, \dots, 2$, $y_1 = y_2$ and $s_1 = s_2$, set $p = p + 1$ if $\pi^{-1}(1) \leq p - 1$, set $\pi = (\pi(1) + 1, \dots, \pi(\pi^{-1}(1) - 1) + 1, 2, 1, \pi(\pi^{-1}(1) + 1) + 1, \dots, \pi(t) + 1)$ in case $s_1 = 1$, and set $\pi = (\pi(1) + 1, \dots, \pi(\pi^{-1}(1) - 1) + 1, 1, 2, \pi(\pi^{-1}(1) + 1) + 1, \dots, \pi(t) + 1)$ in case $s_1 = -1$. Set $t = t + 1$, and go to **Step 1**.

Step 3: Let y^i be the vertex of $D_{v1}(y, \pi, s, p)$ corresponding to the vertex z^- . Set τ_{k+1} equal to the facet of σ_k opposite to the vertex z^- . Consider **Table 1**. If one of the cases BD(j), $j = 1, 2, \dots, 8$, occurs, then τ_{k+1} lies in the boundary of $A(g, \gamma(g))$. Set $k = k + 1$.

1. When one of the cases BD(1), BD(2) and $y_{\pi(1)} = m$, BD(3) and $y_{\pi(i)} = m$, BD(6), or BD(8) occurs, τ_k yields an approximate solution, and the algorithm terminates.

2. When the case both BD(2) and $y_{\pi(1)} = 0$ occurs, set $\kappa = k_1$ and set $g_{k_1} = +1$, $k_j = k_{j+1}$ for $j = 1, 2, \dots, t-2$, $\pi = (\pi(2), \pi(3), \dots, \pi(t))$, $p = p-1$, and $t = t-1$, and go to **Step 4**.
3. When the case both BD(3) and $y_{\pi(i)} = 0$ occurs, set $\kappa = k_1$ and set $g_{k_1} = +1$, $k_j = k_{j+1}$ for $j = 1, 2, \dots, t-2$, $\pi = (\pi(2), \pi(3), \dots, \pi(t))$, and $t = t-1$, and go to **Step 4**.
4. When either the case both BD(4) and either $\pi(i) = 1$ and $s_1 = 1$ or $\pi(i) = 2$ and $s_2 = -1$ or the case both BD(7) and either $\pi(i) = 1$ and $s_1 = 1$ or $\pi(i) = 2$ and $s_2 = -1$ occurs, set $\kappa = k_{t-1}$ and set $g_{k_{t-1}} = -1$, $y_j = y_{j+1}$ and $s_j = s_{j+1}$ for $j = 2, \dots, t-1$, $\pi(j) = \pi(j) - 1$ if $\pi(j) \neq 1$ and 2 for $j < \max\{\pi^{-1}(1), \pi^{-1}(2)\}$, $\pi(\min\{\pi^{-1}(1), \pi^{-1}(2)\}) = 1$ and $\pi(j-1) = \pi(j) - 1$ for $\max\{\pi^{-1}(1), \pi^{-1}(2)\} < j \leq t$, and $t = t-1$, and go to **Step 4**.
5. When the case both BD(5) and that either $\pi(i)$ or $\pi(i+1)$ is equal to one occurs, set $\kappa = k_{t-1}$ and set $g_{k_{t-1}} = -1$, $y_j = y_{j+1}$ and $s_j = s_{j+1}$ for $j = 2, \dots, t-1$, $\pi(j) = \pi(j) - 1$ if $\pi(j) \neq 1$ and 2 for $j < \max\{\pi^{-1}(1), \pi^{-1}(2)\}$, $\pi(\min\{\pi^{-1}(1), \pi^{-1}(2)\}) = 1$ and $\pi(j-1) = \pi(j) - 1$ for $\max\{\pi^{-1}(1), \pi^{-1}(2)\} < j \leq t$, $p = p-1$, and $t = t-1$, and go to **Step 4**.
6. When one of the cases 1) both BD(4) and neither $\pi(i) = 1$ and $s_1 = 1$ nor $\pi(i) = 2$ and $s_2 = -1$, 2) both BD(7) and neither $\pi(i) = 1$ and $s_1 = 1$ nor $\pi(i) = 2$ and $s_2 = -1$, or 3) both BD(5) and that both $\pi(i)$ and $\pi(i+1)$ are unequal to one occurs, set

$$\gamma(g) = (k_1, \dots, k_{t-\pi(i)}, k_{t-\pi(i)-1}, \dots, k_{t-1})$$

if $s_{\pi(i)} = +1$ and set

$$\gamma(g) = (k_1, \dots, k_{t-\pi(i)+1}, k_{t-\pi(i)}, \dots, k_{t-1})$$

if $s_{\pi(i)} = -1$, and go to **Step 1**.

When none of the cases BD(j), $j = 1, 2, \dots, 8$, occurs, set $y = \bar{y}$, $\pi = \bar{\pi}$, $s = s$, and $p = \bar{p}$ according to **Table 1**, and go to **Step 1**.

Step 4: Set σ_{k+1} equal to τ_k and $k = k + 1$, and perform a linear programming step with $-g_\kappa(c(\kappa), 0)^\top$ in the system of linear equations

$$\sum_{j=0}^t \lambda_j \begin{pmatrix} f(z^j) \\ 1 \end{pmatrix} - \sum_{j \notin I^0(g), j \neq \kappa} \mu_j g_j \begin{pmatrix} e^{(j)} \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

If some μ_c becomes zero, then go to **Step 2**; otherwise, some λ_d becomes zero, then set $z^- = z^d$, and go to **Step 3**.

Assuming nondegeneracy in each linear programming step, the algorithm terminates within a finite number of iterations in **Step 2** or the case 1 of **Step 3** with some simplex σ_k . Let z^0, z^1, \dots, z^t be the vertices of σ_k , then $x^1 = \sum_{j=0}^t \lambda_j z^j$ lies in σ_k for $\lambda_j \geq 0$ for all j such that $\sum_{j=0}^t \lambda_j = 1$. Moreover, the vector x^1 can be considered as an approximate complementary point of f on S^n . If the accuracy of approximation is not satisfactory, then the $(2^{n+1} - 2)$ -ray method can be restarted at x^1 for some larger m , in order to improve the accuracy, as has been shown in [5]. Since the D_1 -triangulation is superior to all other well-known triangulations of R^n according to measures of efficiency of triangulations, e.g., see [11] and [12], it is hoped that when using the D_{v1} -triangulation in a simplicial variable dimension algorithm on the unit simplex the cost of computation can be reduced.

References

- [1] Allgower, E.L. and Georg, K.(1980). Simplicial and continuation methods for approximating fixed points and solutions to systems of equations, SIAM Review 22, 28-85.
- [2] Dang, C.(1989).The D_1 -triangulation of R^n for simplicial algorithms for computing solutions of nonlinear equations. Discussion Paper 8928, Center for economic research, Tilburg University. To appear in Mathematics of Operations Research.
- [3] Dang, C. and Talman, A.J.J.(1989). A new triangulation of the unit simplex for computing economic equilibria. Discussion Paper 8935, Center for economic research, Tilburg University. To appear in Methods of Operations Research.

- [4] Doup, T.M.(1988). *Simplicial Algorithms on The Simplotope. Lecture Notes on Economics and Mathematical Systems*, Springer-Verlag, Berlin.
- [5] Doup, T.M., van der Laan, G. and Talman, A.J.J.(1987). The $(2^{n+1} - 2)$ -ray algorithm: a new simplicial algorithm to compute economic equilibria. *Mathematical Programming* 39, 241-252.
- [6] Eaves, B.C.(1984). *A Course in Triangulations for Solving Equations with Deformations. Lecture Notes on Economics and Mathematical Systems*, Springer-Verlag, Berlin.
- [7] Kojima, M. and Yamamoto, Y.(1984). A unified approach to the implementation of several restart fixed point algorithms and a new variable dimension algorithm. *Mathematical Programming* 28, 288-328.
- [8] van der Laan, G. and Talman, A.J.J.(1979). A restart algorithm for computing fixed points without an extra dimension. *Mathematical Programming* 17, 74-84.
- [9] van der Laan, G. and Talman, A.J.J.(1981). A class of simplicial restart fixed point algorithms without an extra dimension. *Mathematical Programming* 20, 33-48.
- [10] Talman, A.J.J. and Yamamoto, Y.(1989). A simplicial algorithm for stationary point problems on polytopes. *Mathematics of Operations Research* 14, 383-399.
- [11] Todd, M.J.(1976). *The Computation of Fixed Points and Applications. Lecture Notes on Economics and Mathematical Systems*, Springer-Verlag, Berlin.
- [12] Todd, M.J.(1976). On triangulations for computing fixed points. *Mathematical Programming* 10, 322-346.
- [13] Wright, A.H.(1981). The octahedral algorithm, a new simplicial fixed point algorithm. *Mathematical Programming* 21, 47-69.
- [14] Yamamoto, Y.(1983). A new variable dimension algorithm for the fixed point problem. *Mathematical Programming* 25, 329-342.

Discussion Paper Series, CentER, Tilburg University, The Netherlands:

(For previous papers please consult previous discussion papers.)

No.	Author(s)	Title
8932	E. van Damme, R. Selten and E. Winter	Alternating Bid Bargaining with a Smallest Money Unit
8933	H. Carlsson and E. van Damme	Global Payoff Uncertainty and Risk Dominance
8934	H. Huizinga	National Tax Policies towards Product- Innovating Multinational Enterprises
8935	C. Dang and D. Talman	A New Triangulation of the Unit Simplex for Computing Economic Equilibria
8936	Th. Nijman and M. Verbeek	The Nonresponse Bias in the Analysis of the Determinants of Total Annual Expenditures of Households Based on Panel Data
8937	A.P. Barten	The Estimation of Mixed Demand Systems
8938	G. Marini	Monetary Shocks and the Nominal Interest Rate
8939	W. Güth and E. van Damme	Equilibrium Selection in the Spence Signaling Game
8940	G. Marini and P. Scaramozzino	Monopolistic Competition, Expected Inflation and Contract Length
8941	J.K. Dagsvik	The Generalized Extreme Value Random Utility Model for Continuous Choice
8942	M.F.J. Steel	Weak Exogeneity in Misspecified Sequential Models
8943	A. Roell	Dual Capacity Trading and the Quality of the Market
8944	C. Hsiao	Identification and Estimation of Dichotomous Latent Variables Models Using Panel Data
8945	R.P. Gilles	Equilibrium in a Pure Exchange Economy with an Arbitrary Communication Structure
8946	W.B. MacLeod and J.M. Malcomson	Efficient Specific Investments, Incomplete Contracts, and the Role of Market Alterna- tives
8947	A. van Soest and A. Kapteyn	The Impact of Minimum Wage Regulations on Employment and the Wage Rate Distribution
8948	P. Kooreman and B. Melenberg	Maximum Score Estimation in the Ordered Response Model

No.	Author(s)	Title
8949	C. Dang	The D_3 -Triangulation for Simplicial Deformation Algorithms for Computing Solutions of Nonlinear Equations
8950	M. Cripps	Dealer Behaviour and Price Volatility in Asset Markets
8951	T. Wansbeek and A. Kapteyn	Simple Estimators for Dynamic Panel Data Models with Errors in Variables
8952	Y. Dai, G. van der Laan, D. Talman and Y. Yamamoto	A Simplicial Algorithm for the Nonlinear Stationary Point Problem on an Unbounded Polyhedron
8953	F. van der Ploeg	Risk Aversion, Intertemporal Substitution and Consumption: The CARA-LQ Problem
8954	A. Kapteyn, S. van de Geer, H. van de Stadt and T. Wansbeek	Interdependent Preferences: An Econometric Analysis
8955	L. Zou	Ownership Structure and Efficiency: An Incentive Mechanism Approach
8956	P. Kooreman and A. Kapteyn	On the Empirical Implementation of Some Game Theoretic Models of Household Labor Supply
8957	E. van Damme	Signaling and Forward Induction in a Market Entry Context
9001	A. van Soest, P. Kooreman and A. Kapteyn	Coherency and Regularity of Demand Systems with Equality and Inequality Constraints
9002	J.R. Magnus and B. Pesaran	Forecasting, Misspecification and Unit Roots: The Case of AR(1) Versus ARMA(1,1)
9003	J. Driffill and C. Schultz	Wage Setting and Stabilization Policy in a Game with Renegotiation
9004	M. McAleer, M.H. Pesaran and A. Bera	Alternative Approaches to Testing Non-Nested Models with Autocorrelated Disturbances: An Application to Models of U.S. Unemployment
9005	Th. ten Raa and M.F.J. Steel	A Stochastic Analysis of an Input-Output Model: Comment
9006	M. McAleer and C.R. McKenzie	Keynesian and New Classical Models of Unemployment Revisited
9007	J. Osiewalski and M.F.J. Steel	Semi-Conjugate Prior Densities in Multivariate t Regression Models

No.	Author(s)	Title
9007	J. Osiewalski and M.F.J. Steel	Semi-Conjugate Prior Densities in Multi- variate t Regression Models
9008	G.W. Imbens	Duration Models with Time-Varying Coefficients
9009	G.W. Imbens	An Efficient Method of Moments Estimator for Discrete Choice Models with Choice-Based Sampling
9010	P. Deschamps	Expectations and Intertemporal Separability in an Empirical Model of Consumption and Investment under Uncertainty
9011	W. Güth and E. van Damme	Gorby Games - A Game Theoretic Analysis of Disarmament Campaigns and the Defense Efficiency-Hypothesis
9012	A. Horsley and A. Wrobel	The Existence of an Equilibrium Density for Marginal Cost Prices, and the Solution to the Shifting-Peak Problem
9013	A. Horsley and A. Wrobel	The Closedness of the Free-Disposal Hull of a Production Set
9014	A. Horsley and A. Wrobel	The Continuity of the Equilibrium Price Density: The Case of Symmetric Joint Costs, and a Solution to the Shifting-Pattern Problem
9015	A. van den Elzen, G. van der Laan and D. Talman	An Adjustment Process for an Exchange Economy with Linear Production Technologies
9016	P. Deschamps	On Fractional Demand Systems and Budget Share Positivity
9017	B.J. Christensen and N.M. Kiefer	The Exact Likelihood Function for an Empirical Job Search Model
9018	M. Verbeek and Th. Nijman	Testing for Selectivity Bias in Panel Data Models
9019	J.R. Magnus and B. Pesaran	Evaluation of Moments of Ratios of Quadratic Forms in Normal Variables and Related Statistics
9020	A. Robson	Status, the Distribution of Wealth, Social and Private Attitudes to Risk
9021	J.R. Magnus and B. Pesaran	Evaluation of Moments of Quadratic Forms in Normal Variables

No.	Author(s)	Title
9022	K. Kamiya and D. Talman	Linear Stationary Point Problems
9023	W. Emons	Good Times, Bad Times, and Vertical Upstream Integration
9024	C. Dang	The D_2 -Triangulation for Simplicial Homotopy Algorithms for Computing Solutions of Nonlinear Equations
9025	K. Kamiya and D. Talman	Variable Dimension Simplicial Algorithm for Balanced Games
9026	P. Skott	Efficiency Wages, Mark-Up Pricing and Effective Demand
9027	C. Dang and D. Talman	The D_1 -Triangulation in Simplicial Variable Dimension Algorithms for Computing Solutions of Nonlinear Equations
9028	J. Bai, A.J. Jakeman and M. McAleer	Discrimination Between Nested Two- and Three- Parameter Distributions: An Application to Models of Air Pollution
9029	Th. van de Klundert	Crowding out and the Wealth of Nations
9030	Th. van de Klundert and R. Gradus	Optimal Government Debt under Distortionary Taxation
9031	A. Weber	The Credibility of Monetary Target Announce- ments: An Empirical Evaluation
9032	J. Osiewalski and M. Steel	Robust Bayesian Inference in Elliptical Regression Models
9033	C. R. Wichers	The Linear-Algebraic Structure of Least Squares
9034	C. de Vries	On the Relation between GARCH and Stable Processes
9035	M.R. Baye, D.W. Jansen and Q. Li	Aggregation and the "Random Objective" Justification for Disturbances in Complete Demand Systems
9036	J. Driffill	The Term Structure of Interest Rates: Structural Stability and Macroeconomic Policy Changes in the UK
9037	F. van der Ploeg	Budgetary Aspects of Economic and Monetary Integration in Europe

No.	Author(s)	Title
9038	A. Robson	Existence of Nash Equilibrium in Mixed Strategies for Games where Payoffs Need not Be Continuous in Pure Strategies
9039	A. Robson	An "Informationally Robust Equilibrium" for Two-Person Nonzero-Sum Games
9040	M.R. Baye, G. Tian and J. Zhou	The Existence of Pure-Strategy Nash Equilibrium in Games with Payoffs that are not Quasiconcave
9041	M. Burnovsky and I. Zang	"Costless" Indirect Regulation of Monopolies with Substantial Entry Cost
9042	P.J. Deschamps	Joint Tests for Regularity and Autocorrelation in Allocation Systems
9043	S. Chib, J. Osiewalski and M. Steel	Posterior Inference on the Degrees of Freedom Parameter in Multivariate-t Regression Models
9044	H.A. Keuzenkamp	The Probability Approach in Economic Methodology: On the Relation between Haavelmo's Legacy and the Methodology of Economics
9045	I.M. Bomze and E.F.C. van Damme	A Dynamical Characterization of Evolutionarily Stable States
9046	E. van Damme	On Dominance Solvable Games and Equilibrium Selection Theories
9047	J. Driffill	Changes in Regime and the Term Structure: A Note
9048	A.J.J. Talman	General Equilibrium Programming
9049	H.A. Keuzenkamp and F. van der Ploeg	Saving, Investment, Government Finance and the Current Account: The Dutch Experience
9050	C. Dang and A.J.J. Talman	The D_1 -Triangulation in Simplicial Variable Dimension Algorithms on the Unit Simplex for Computing Fixed Points

P.O. BOX 90153. 5000 LE TILBURG. THE NETHERLANDS

Bibliotheek K. U. Brabant



17 000 01117575 0